

Gimoti®

(metoclopramide)
nasal spray

Instructions for Use



What is Gimoti® (metoclopramide) nasal spray?

GIMOTI is a prescription medicine used 30 minutes before meals and at bedtime for 2 to 8 weeks to relieve symptoms in adults with diabetes who have irregular or slow stomach emptying. Avoid treatment with metoclopramide (all dosage forms and routes of administration) for longer than 12 weeks.

GIMOTI is not recommended for use in children under age 18.


Read these Instructions for Use before you start using GIMOTI and each time you get a refill. This information does not take the place of talking to your healthcare professional about your medical condition or your treatment.



Important Information

- GIMOTI is for use in your nose only.
- Each GIMOTI bottle has enough medicine for 4 weeks of treatment when taken 4 times each day (recommended dosage).
- One dose is 1 spray in 1 nostril.
- Your GIMOTI bottle must be primed
 - before you use it for the first time
 - when the nasal spray has not been used for 2 weeks
- Store the GIMOTI bottle at 68°F to 77°F (20°C to 25°C).
- Throw away the GIMOTI bottle 4 weeks after opening even if the bottle contains unused medicine.
- Keep out of the reach of children.


Parts of your GIMOTI bottle



Steps to use GIMOTI

Step 1 » Uncap the GIMOTI bottle

- Remove the cap from the spray nozzle by pulling straight up.




See next page for continued instructions.

Please see complete Important Safety Information attached, including Boxed Warning.

Step 2 » Prime the GIMOTI bottle

- Your GIMOTI bottle must be primed
 - before you use it for the first time
 - when the nasal spray has not been used for 2 weeks or longer
- Remove the safety clip from the spray pump.
- Place your index finger and middle finger on each side of the finger flange and your thumb on the bottom of the glass bottle.
- Turn the spray nozzle away from your face, keeping the bottle upright. **Do not** spray into your eyes.
- Press down firmly and release 10 times on the finger flange until a spray appears from the spray nozzle. You may not see a spray the first few times you press down. After pressing down and releasing a spray 10 times, the GIMOTI bottle will be primed and ready to use.

If you are not able to press and release 10 sprays from the GIMOTI bottle, call your healthcare professional or pharmacist.



The GIMOTI bottle is now ready for use.

Step 3 » Use the GIMOTI bottle


- Place the spray nozzle tip under one of your nostrils. Tilt your head slightly forward so the tip of the spray nozzle is aimed away from the center of your nose.
- Close the other nostril with your other index finger. Move the spray pump up so the tip of the nozzle is in your nostril.
- Hold the bottle upright and press down firmly on the finger flange until it stops moving to deliver a full dose.
- Release the finger flange and breathe in gently through the open nostril.
- Remove the spray pump nozzle tip from your nostril and breathe out slowly through your mouth.
- After use, wipe the spray nozzle with a clean tissue.

Replace the cap on the spray nozzle by pushing straight down. Place the safety clip back onto the spray pump.



Step 4 » Clean the GIMOTI spray pump nozzle

- If the spray pump nozzle becomes clogged, remove it for cleaning by grasping the base of the spray nozzle and pulling up.
- Soak the spray nozzle in warm water and rinse. Do not try to unblock the spray nozzle by inserting a pin or other sharp object because this will damage the spray nozzle.
- Dry the spray nozzle at room temperature. When the spray nozzle is dry, place it back on the GIMOTI bottle.



Disposal instructions: If there is no take-back program available in your area, throw away GIMOTI bottles in the household trash.

Please see complete Important Safety Information attached, including Boxed Warning.

IMPORTANT SAFETY INFORMATION—PATIENT

Gimoti® (metoclopramide) nasal spray

What is Gimoti® (metoclopramide) nasal spray?

GIMOTI is a prescription medicine used 30 minutes before meals and at bedtime to relieve symptoms of slow stomach emptying in adults with diabetes.

GIMOTI is not recommended for use in children under age 18.

Important facts about GIMOTI

This is a summary of important information you need to know about GIMOTI. This information does not take the place of talking with your healthcare professional about your medical condition or your treatment.

What is the most important information I should know about GIMOTI?

GIMOTI can cause serious side effects, including: TARDIVE DYSKINESIA: Abnormal muscle movements, mostly of the face or tongue muscles. You cannot control these movements, and they may not go away even after stopping GIMOTI. Your chances of getting tardive dyskinesia increase:

- **The longer you take metoclopramide and the more metoclopramide you take. You should not take GIMOTI for more than 12 weeks. If you take GIMOTI for more than 12 weeks, it is important to monitor for signs and symptoms of TD.**
- **If you are older, especially if you are an older woman (e.g., age 65 years and older)**
- **If you have diabetes**

Stop taking GIMOTI and call your healthcare professional right away if you get movements you cannot stop or control, such as lip smacking, chewing, or puckering up your mouth; frowning or scowling; sticking out your tongue; blinking and moving your eyes; shaking of your arms and legs.

Do not use GIMOTI if you:

- Have a history of tardive dyskinesia or have a problem controlling your muscles and movements after taking GIMOTI or a medicine that works like GIMOTI
- Have stomach or intestinal problems that could get worse with GIMOTI, such as bleeding, blockage, or a tear in the stomach or bowel wall
- Have a type of tumor that can cause high blood pressure, such as pheochromocytoma
- Have epilepsy (seizures)
- Are allergic to metoclopramide. Stop taking GIMOTI right away and get emergency help if you have any of these symptoms:
 - swelling of your tongue, throat, lips, eyes, or face
 - trouble swallowing or breathing
 - skin rash, hives, sores in your mouth, or skin blisters

Before starting GIMOTI, tell your healthcare professional about all your medical conditions, especially if you have:

- Problems controlling your muscle movements after taking any medicine
- Parkinson's disease
- Pheochromocytoma
- Kidney or liver disease
- Depression or mental illness
- High blood pressure

Also tell your healthcare professional if you are pregnant or plan to become pregnant, are breastfeeding or plan to breastfeed, or drink alcohol.

Tell your healthcare professional about all the medicines you take, including prescription drugs and over-the-counter medicines, vitamins, and herbal supplements.

Speak with your healthcare professional before you start or stop any other medicines.

Especially tell your healthcare professional if you take:

- Another medicine that contains metoclopramide, such as REGLAN® tablets
- Medicine for Parkinson's disease
- Blood pressure medicine
- Medicine for depression, especially a monoamine oxidase inhibitor (MAOI)
- Antipsychotic medicine used to treat mental illness, such as schizophrenia
- Insulin
- Medicines that can make you sleepy, such as anxiety medicines, sleep medicines, and narcotics

What should I avoid while taking GIMOTI?

- Do not drink alcohol while taking GIMOTI.
- GIMOTI may cause sleepiness or dizziness. Do not drive, operate machinery, or do potentially dangerous activities until you know how GIMOTI affects you.

What are other possible side effects of GIMOTI?

- Other changes in muscle control and movement, such as:
 - uncontrolled spasms of your face and neck muscles, or muscles of your body, arms, and legs (dystonia)
 - parkinsonism – slight shaking, body stiffness, and trouble moving or keeping your balance
 - being unable to sit still or feeling that you need to move your hands, feet, or body (akathisia)
- Neuroleptic Malignant Syndrome (NMS) – a very rare but very serious condition. NMS can lead to death and must be treated in a hospital.
- Depression, thoughts about suicide, and suicide
- High blood pressure
- Too much body water
- Increased prolactin

Call your healthcare professional and get medical help right away if you:

- Feel depressed or have thoughts about hurting or killing yourself
- Have high fever, stiff muscles, problems thinking, very fast or uneven heartbeat, and/or increased sweating
- Have muscle movements that you cannot stop or control
- Have muscle movements that are new or unusual

The most common side effects of GIMOTI include:

- Unpleasant taste after dosing
- Headache
- Tiredness

These are not all the possible side effects of GIMOTI. Ask your healthcare professional for medical advice about side effects.

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.FDA.gov/medwatch or call 1-800-FDA-1088.

This information should not take the place of you talking with your doctor or healthcare professional. If you have any questions about your condition, or if you would like more information about GIMOTI, talk to your doctor or pharmacist. Only you and your healthcare professional can decide if GIMOTI is right for you.

HIGHLIGHTS OF PRESCRIBING INFORMATION
These highlights do not include all of the information needed to use GIMOTI safely and effectively. See full prescribing information for GIMOTI.

GIMOTI® (metoclopramide) nasal spray
Initial U.S. Approval: 1979

WARNING: TARDIVE DYSKINESIA	
See full prescribing information for complete boxed warning.	
Metoclopramide, including GIMOTI, can cause tardive dyskinesia (TD), a potentially irreversible serious movement disorder. In patients treated with metoclopramide, including GIMOTI, the risk of developing TD increases with duration of treatment and total cumulative dosage. (5.1) GIMOTI is contraindicated in patients with a history of TD. (4) Use GIMOTI for the shortest duration of treatment and periodically reassess the need for continued treatment. (2.2, 5.1) Immediately discontinue GIMOTI in patients who develop signs or symptoms of TD. (5.1) Avoid a total duration of treatment with metoclopramide products, including GIMOTI, for longer than 12 weeks. If longer-term use is unavoidable, routinely monitor for signs and symptoms of TD. (5.1)	

RECENT MAJOR CHANGES	
Boxed Warning	2/2026
Dosage and Administration, Recommended Dosage (2.2)	2/2026
Warnings and Precautions, Tardive Dyskinesia (5.1)	2/2026
Warnings and Precautions Other Extrapyramidal Symptoms (5.2)	2/2026
INDICATIONS AND USAGE	
GIMOTI is a dopamine-2 (D ₂) antagonist indicated for the relief of symptoms in adults with acute and recurrent diabetic gastroparesis. (1)	
Limitations of Use:	
GIMOTI is not recommended for use in:	
- pediatric patients due to the risk of tardive dyskinesia (TD) and other extrapyramidal symptoms as well as the risk of methemoglobinemia in neonates. (1, 8.4) - moderate or severe hepatic impairment (Child-Pugh B or C), moderate or severe renal impairment (creatinine clearance less than 60 mL/minute), and patients concurrently using strong CYP2D6 inhibitors due to the risk of increased drug exposure and adverse reactions. (1, 5.9, 7.1)	
DOSAGE AND ADMINISTRATION	

Administration
• See the full prescribing information for complete information on administration. (2.1)

Boxed Warning	
2/2026	
Dosage and Administration, Recommended Dosage (2.2)	
2/2026	
Warnings and Precautions, Tardive Dyskinesia (5.1)	
2/2026	
Warnings and Precautions Other Extrapyramidal Symptoms (5.2)	
2/2026	

INDICATIONS AND USAGE
GIMOTI is a dopamine-2 (D₂) antagonist indicated for the relief of symptoms in adults with acute and recurrent diabetic gastroparesis. (1)

Limitations of Use:
GIMOTI is not recommended for use in:

- pediatric patients due to the risk of tardive dyskinesia (TD) and other extrapyramidal symptoms as well as the risk of methemoglobinemia in neonates. (1, 8.4)
- moderate or severe hepatic impairment (Child-Pugh B or C), moderate or severe renal impairment (creatinine clearance less than 60 mL/minute), and patients concurrently using strong CYP2D6 inhibitors due to the risk of increased drug exposure and adverse reactions. (1, 5.9, 7.1)

DOSAGE AND ADMINISTRATION
GIMOTI is a dopamine-2 (D₂) antagonist indicated for the relief of symptoms in adults with acute and recurrent diabetic gastroparesis. (1)

Limitations of Use:
GIMOTI is not recommended for use in:

- pediatric patients due to the risk of tardive dyskinesia (TD) and other extrapyramidal symptoms as well as the risk of methemoglobinemia in neonates. (1, 8.4)
- moderate or severe hepatic impairment (Child-Pugh B or C), moderate or severe renal impairment (creatinine clearance less than 60 mL/minute), and patients concurrently using strong CYP2D6 inhibitors due to the risk of increased drug exposure and adverse reactions. (1, 5.9, 7.1)

DOSAGE AND ADMINISTRATION
GIMOTI is a dopamine-2 (D₂) antagonist indicated for the relief of symptoms in adults with acute and recurrent diabetic gastroparesis. (1)

Limitations of Use:
GIMOTI is not recommended for use in:

- pediatric patients due to the risk of tardive dyskinesia (TD) and other extrapyramidal symptoms as well as the risk of methemoglobinemia in neonates. (1, 8.4)
- moderate or severe hepatic impairment (Child-Pugh B or C), moderate or severe renal impairment (creatinine clearance less than 60 mL/minute), and patients concurrently using strong CYP2D6 inhibitors due to the risk of increased drug exposure and adverse reactions. (1, 5.9, 7.1)

DOSAGE AND ADMINISTRATION
GIMOTI is a dopamine-2 (D₂) antagonist indicated for the relief of symptoms in adults with acute and recurrent diabetic gastroparesis. (1)

Limitations of Use:
GIMOTI is not recommended for use in:

- pediatric patients due to the risk of tardive dyskinesia (TD) and other extrapyramidal symptoms as well as the risk of methemoglobinemia in neonates. (1, 8.4)
- moderate or severe hepatic impairment (Child-Pugh B or C), moderate or severe renal impairment (creatinine clearance less than 60 mL/minute), and patients concurrently using strong CYP2D6 inhibitors due to the risk of increased drug exposure and adverse reactions. (1, 5.9, 7.1)

DOSAGE AND ADMINISTRATION
GIMOTI is a dopamine-2 (D₂) antagonist indicated for the relief of symptoms in adults with acute and recurrent diabetic gastroparesis. (1)

Limitations of Use:
GIMOTI is not recommended for use in:

- pediatric patients due to the risk of tardive dyskinesia (TD) and other extrapyramidal symptoms as well as the risk of methemoglobinemia in neonates. (1, 8.4)
- moderate or severe hepatic impairment (Child-Pugh B or C), moderate or severe renal impairment (creatinine clearance less than 60 mL/minute), and patients concurrently using strong CYP2D6 inhibitors due to the risk of increased drug exposure and adverse reactions. (1, 5.9, 7.1)

DOSAGE AND ADMINISTRATION
GIMOTI is a dopamine-2 (D₂) antagonist indicated for the relief of symptoms in adults with acute and recurrent diabetic gastroparesis. (1)

Limitations of Use:
GIMOTI is not recommended for use in:

- pediatric patients due to the risk of tardive dyskinesia (TD) and other extrapyramidal symptoms as well as the risk of methemoglobinemia in neonates. (1, 8.4)
- moderate or severe hepatic impairment (Child-Pugh B or C), moderate or severe renal impairment (creatinine clearance less than 60 mL/minute), and patients concurrently using strong CYP2D6 inhibitors due to the risk of increased drug exposure and adverse reactions. (1, 5.9, 7.1)

DOSAGE AND ADMINISTRATION
GIMOTI is a dopamine-2 (D₂) antagonist indicated for the relief of symptoms in adults with acute and recurrent diabetic gastroparesis. (1)

Limitations of Use:
GIMOTI is not recommended for use in:

- pediatric patients due to the risk of tardive dyskinesia (TD) and other extrapyramidal symptoms as well as the risk of methemoglobinemia in neonates. (1, 8.4)
- moderate or severe hepatic impairment (Child-Pugh B or C), moderate or severe renal impairment (creatinine clearance less than 60 mL/minute), and patients concurrently using strong CYP2D6 inhibitors due to the risk of increased drug exposure and adverse reactions. (1, 5.9, 7.1)

DOSAGE AND ADMINISTRATION
GIMOTI is a dopamine-2 (D₂) antagonist indicated for the relief of symptoms in adults with acute and recurrent diabetic gastroparesis. (1)

Limitations of Use:
GIMOTI is not recommended for use in:

- pediatric patients due to the risk of tardive dyskinesia (TD) and other extrapyramidal symptoms as well as the risk of methemoglobinemia in neonates. (1, 8.4)
- moderate or severe hepatic impairment (Child-Pugh B or C), moderate or severe renal impairment (creatinine clearance less than 60 mL/minute), and patients concurrently using strong CYP2D6 inhibitors due to the risk of increased drug exposure and adverse reactions. (1, 5.9, 7.1)

DOSAGE AND ADMINISTRATION
GIMOTI is a dopamine-2 (D₂) antagonist indicated for the relief of symptoms in adults with acute and recurrent diabetic gastroparesis. (1)

Limitations of Use:
GIMOTI is not recommended for use in:

- pediatric patients due to the risk of tardive dyskinesia (TD) and other extrapyramidal symptoms as well as the risk of methemoglobinemia in neonates. (1, 8.4)
- moderate or severe hepatic impairment (Child-Pugh B or C), moderate or severe renal impairment (creatinine clearance less than 60 mL/minute), and patients concurrently using strong CYP2D6 inhibitors due to the risk of increased drug exposure and adverse reactions. (1, 5.9, 7.1)

DOSAGE AND ADMINISTRATION
GIMOTI is a dopamine-2 (D₂) antagonist indicated for the relief of symptoms in adults with acute and recurrent diabetic gastroparesis. (1)

RECOMMENDED DOSAGE
Adults less than 65 years of age: The recommended dosage is 1 spray (15 mg) in one nostril, 30 minutes before each meal and at bedtime (maximum of 4 sprays daily). (2,2)

Adults 65 years of age and older: GIMOTI is not recommended in geriatric patients as initial therapy. Geriatric patients receiving an alternative metoclopramide product at a stable dosage of 10 mg four times daily can be switched to GIMOTI 1 spray (15 mg) in one nostril, 30 minutes before each meal and at bedtime (maximum four times daily). (2.2, 8.5)

Nasal Spray: 15 mg metoclopramide in each 70 microliter spray. (3)

CONTRAINDICATIONS
History of TD or dystonic reaction to metoclopramide (4)

When stimulation of gastrointestinal motility might be dangerous (e.g., in the presence of gastrointestinal hemorrhage, mechanical obstruction, or perforation).

In patients with pheochromocytoma or other catecholamine-releasing paragangliomas. Metoclopramide may cause a hypertensive/pheochromocytoma crisis, probably due to release of catecholamines from the tumor [see Warnings and Precautions (5.5)].

In patients with epilepsy. Metoclopramide may increase the frequency and severity of seizures [see Adverse Reactions (6)].

In patients with hypersensitivity to metoclopramide. Reactions have included laryngeal and glossal angioedema and bronchospasm [see Adverse Reactions (6)].

WARNINGS AND PRECAUTIONS
Tardive dyskinesia (TD), other extrapyramidal symptoms (EPS), and neuroleptic malignant syndrome (NMS): Avoid concomitant use of other drugs known to cause TD/EPS/NMS and avoid use in patients with Parkinson's disease. If symptoms occur, discontinue GIMOTI and seek immediate medical attention. (5.1, 5.2, 5.3, 7.1, 7.2)

Depression and suicidal ideation/suicide: Avoid use. (5.4)

ADVERSE REACTIONS
Most common adverse reactions (≥5%) are: dysgeusia, headache, and fatigue. (6)

DRUG INTERACTIONS
Antipsychotics: Potential for additive effects, including TD, EPS, and NMS; avoid concomitant use. (7.1)

Central nervous system (CNS) depressants: Increased risk of CNS depression. Avoid concomitant use and monitor for adverse reactions. (7.1)

Monoamine oxidase (MAO) inhibitors: Increased risk of hypertension; avoid concomitant use. (5.5, 7.1)

Additional drug interactions: See Full Prescribing Information. (7.1, 7.2)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 4/2026

8 USE IN SPECIFIC POPULATIONS
8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Renal Impairment

8.7 Hepatic Impairment

8.8 NADH-Cytochrome b₅ Reductase Deficiency

8.9 CYP2D6 Poor Metabolizers

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

Storage
Discard GIMOTI 4 weeks after opening even if the bottle contains unused drug.

2.2 Recommended Dosage
Adults Less Than 65 Years of Age

The recommended dosage of GIMOTI for the treatment of acute and recurrent diabetic gastroparesis in adults is 1 spray (15 mg) in one nostril, 30 minutes before each meal and at bedtime (maximum of four times daily).

Adults 65 Years of Age and Older
Elderly patients may be more sensitive to the adverse effects of metoclopramide and require a lower starting dosage [see Warnings and Precautions (5.1)]. GIMOTI is not recommended in geriatric patients as initial therapy.

Geriatric patients receiving an alternative metoclopramide product at a stable dosage of 10 mg four times daily can be switched to GIMOTI 1 spray (15 mg) in one nostril, 30 minutes before each meal and at bedtime (maximum four times daily).

Treatment Duration
Use GIMOTI for the shortest duration of treatment and periodically reassess the need for continued treatment.

Avoid treatment with metoclopramide, including GIMOTI, for longer than 12 weeks. If longer-term use is unavoidable, routinely monitor for signs and symptoms of TD [see Warnings and Precautions (5.1)].

3 DOSAGE FORMS AND STRENGTHS
Nasal Spray: 15 mg of metoclopramide in each 70 microliter spray. GIMOTI is an aqueous solution supplied in an amber glass bottle fitted with a metered spray pump attachment.

4 CONTRAINDICATIONS
GIMOTI is contraindicated:

In patients with a history of tardive dyskinesia (TD) or a dystonic reaction to metoclopramide [see Warnings and Precautions (5.1, 5.2)].

When stimulation of gastrointestinal motility might be dangerous (e.g., in the presence of gastrointestinal hemorrhage, mechanical obstruction, or perforation).

In patients with pheochromocytoma or other catecholamine-releasing paragangliomas. Metoclopramide may cause a hypertensive/pheochromocytoma crisis, probably due to release of catecholamines from the tumor [see Warnings and Precautions (5.5)].

In patients with epilepsy. Metoclopramide may increase the frequency and severity of seizures [see Adverse Reactions (6)].

In patients with hypersensitivity to metoclopramide. Reactions have included laryngeal and glossal angioedema and bronchospasm [see Adverse Reactions (6)].

5 WARNINGS AND PRECAUTIONS
5.1 Tardive Dyskinesia

Metoclopramide, including GIMOTI, can cause tardive dyskinesia (TD), a syndrome of potentially irreversible and disfiguring involuntary movements of the face or tongue, and sometimes of the trunk and/or extremities. Metoclopramide, including GIMOTI, may also suppress, or partially suppress, the signs of TD, and may delay the diagnosis of TD because it may mask the underlying disease process. The effect of this symptomatic suppression upon the long-term course of TD is unknown. TD may remit, partially or completely, if GIMOTI treatment is discontinued.

In patients treated with metoclopramide, including GIMOTI, the risk of developing TD and the likelihood that TD will become irreversible increases with duration of treatment and total cumulative dosage. Additionally, the risk of developing TD is increased in elderly patients, especially in elderly women [see Use in Specific Populations (8.5)], and in patients with diabetes mellitus.

Prevention, Mitigation, and Monitoring for TD

GIMOTI is contraindicated in patients with a history of TD.

Avoid use of GIMOTI in patients receiving concomitant antipsychotics due to the potential additive effects of TD [see Drug Interactions (7.1)].

GIMOTI is not recommended in geriatric patients as initial therapy [see Dosage and Administration (2.2)].

Use GIMOTI for the shortest duration of treatment and periodically reassess the need for continued treatment.

Immediately discontinue GIMOTI immediately in patients who develop signs and symptoms of TD.

Avoid a total duration of treatment with metoclopramide products, including GIMOTI, for longer than 12 weeks. If longer-term use is unavoidable, routinely monitor for signs and symptoms of TD.

If patients have continued TD symptoms, consider TD treatment.

5.2 Other Extrapyramidal Symptoms

In addition to TD, metoclopramide may cause other extrapyramidal symptoms (EPS), parkinsonian symptoms, and motor restlessness. Advise patients to seek immediate medical attention if such symptoms occur and to discontinue GIMOTI.

Extrapyramidal symptoms (EPS), such as acute dystonic reactions, occurred in patients treated with oral metoclopramide dosages of 30 mg to 40 mg daily. Such reactions occurred more frequently in adults less than 30 years of age and at higher than recommended dosages. EPS occurred more frequently in pediatric patients compared to adults (GIMOTI is not approved for use in pediatric patients). Symptoms can occur in the first 24 to 48 hours after starting metoclopramide. Symptoms included involuntary movements of limbs and facial grimacing, torticollis, oculogyric crisis, rhythmic protrusion of tongue, bulbar type of speech, trismus, or dystonic reactions resembling tetanus. Rarely, dystonic reactions were present as stridor and dyspnea, possibly due to laryngospasm. Diphenhydramine hydrochloride or benztropine mesylate may be used to treat these adverse reactions. Avoid GIMOTI in patients receiving other drugs that can cause EPS (e.g., antipsychotics).

Parkinsonian symptoms (bradykinesia, tremor, cogwheel rigidity, mask-like facies) have occurred after starting metoclopramide, more commonly within the first 6 months, but also after longer periods. Symptoms generally have subsided within 2 to 3 months after discontinuation of metoclopramide. Avoid GIMOTI in patients with Parkinson's disease and other patients being treated with antiparkinsonian drugs due to potential exacerbation of symptoms. If treatment is unavoidable, use GIMOTI for the shortest duration of treatment and periodically reassess the need for continued treatment. Routinely monitor for signs and symptoms of Parkinson's disease [see Dosage and Administration (2.2)].

Motor restlessness (akathisia) has developed and consisted of feelings of anxiety, agitation, jitteriness, and insomnia, as well as inability to sit still, pacing, and foot tapping. Discontinue GIMOTI if these symptoms develop.

5.3 Neuroleptic Malignant Syndrome

Metoclopramide may cause a potentially fatal symptom complex called neuroleptic malignant syndrome (NMS). NMS has been reported in association with metoclopramide overdose and concomitant treatment with another drug associated with NMS. Avoid GIMOTI in patients receiving other drugs associated with NMS, including typical and atypical antipsychotics.

Clinical manifestations of NMS include hyperpyrexia, muscle rigidity, altered mental status, and manifestations of autonomic instability (irregular pulse or blood pressure, tachycardia, diaphoresis, and cardiac arrhythmias). Additional signs may include elevated creatine phosphokinase, myoglobinuria (rhabdomyolysis), and acute renal failure. Patients with such symptoms should be evaluated immediately.

In the diagnostic evaluation, consider the presence of other serious medical conditions (e.g., pneumonia, systemic infection) and untreated or inadequately treated extrapyramidal signs and symptoms. Other important considerations in the differential diagnosis include central anticholinergic toxicity, heat stroke, malignant hyperthermia, drug fever, serotonin syndrome, and primary central nervous system pathology.

Management of NMS includes:

Immediate discontinuation of GIMOTI and other drugs not essential to concurrent therapy [see Drug Interactions (7.1)].

Intensive symptomatic treatment and medical monitoring.

Treatment of any concomitant serious medical problems for which specific treatments are available.

7 DRUG INTERACTIONS
7.1 Effects of Other Drugs on Metoclopramide

Table 1 displays the effects of other drugs on metoclopramide.

Table 1. Effects of Other Drugs on Metoclopramide

Antipsychotics

Clinical Impact
Potential for additive effects, including increased frequency and severity of tardive dyskinesia (TD), other extrapyramidal symptoms (EPS), and neuroleptic malignant syndrome (NMS).

Intervention
Avoid concomitant use [see Warnings and Precautions (5.1, 5.2, 5.3)].

Strong CYP2D6 Inhibitors, not Included in Antipsychotic Category Above

Clinical Impact
Increased plasma concentrations of metoclopramide; risk of exacerbation of extrapyramidal symptoms [see Clinical Pharmacology (12.3)].

Intervention
Use of GIMOTI is not recommended [see Warnings and Precautions (5.9)].

Examples
quinidine, bupropion, fluoxetine, and paroxetine

Monoamine Oxidase Inhibitors

Clinical Impact
Increased risk of hypertension [see Warnings and Precautions (5.5)].

Intervention
Avoid concomitant use.

Central Nervous System (CNS) Depressants

Clinical Impact
Increased risk of CNS depression [see Warnings and Precautions (5.8)].

Intervention
Avoid GIMOTI or the interacting drug, depending on the importance of the drug to the patient.

Examples
alcohol, sedatives, hypnotics, opiates, and anxiolytics

Drugs that Impair Gastrointestinal Motility

Clinical Impact
Potential for decreased effect of metoclopramide due to opposing effects on gastrointestinal motility.

Intervention
Monitor for reduced effect on gastrointestinal motility.

Examples
antiperistaltic antidiarrheal drugs, anticholinergic drugs, and opiates

Dopaminergic Agonists and Other Drugs that Increase Dopamine Concentrations

Clinical Impact
Decreased therapeutic effect of metoclopramide due to opposing effects on dopamine.

Intervention
Monitor for reduced therapeutic effect.

Examples
apomorphine, bromocriptine, cabergoline, levodopa, pramipexole, ropinirole, and rotigotine

7.2 Effects of Metoclopramide on Other Drugs

Table 2 displays the effects of metoclopramide on other drugs.

Table 2. Effects of Metoclopramide on Other Drugs

Dopaminergic Agonists and Drugs Increasing Dopamine Concentrations

Clinical Impact
Opposing effects of metoclopramide and the interacting drug on dopamine. Potential exacerbation of symptoms (e.g., parkinsonian symptoms).

Intervention
Avoid concomitant use [see Warnings and Precautions (5.2)].

Examples
Apomorphine, bromocriptine, cabergoline, levodopa, pramipexole, ropinirole, rotigotine

Succinylcholine, Mivacurium

Clinical Impact
Metoclopramide inhibits plasma cholinesterase leading to enhanced neuromuscular blockade.

Intervention
Monitor for signs and symptoms of prolonged neuromuscular blockade

Drugs with Absorption Altered due to Increased Gastrointestinal Motility

Clinical Impact
The effect of metoclopramide on other drugs is variable. Increased gastrointestinal (GI) motility by metoclopramide may impact absorption of other drugs leading to decreased or increased drug exposure.

Intervention
Drugs with Decreased Absorption (e.g., digoxin, atovaquone, posaconazole oral suspension*, fosfomycin): Monitor for reduced therapeutic effect of the interacting drug. For digoxin, monitor therapeutic drug concentrations and increase the digoxin dose as needed (see prescribing information for digoxin).

Drugs with Increased Absorption (e.g., sirolimus, tacrolimus, cyclosporine): Monitor therapeutic drug concentrations and adjust the dose as needed. See prescribing information for the interacting drug.

Insulin

Clinical Impact
Increased GI motility by metoclopramide may increase delivery of food to the intestines and increase blood glucose.

Intervention
Monitor blood glucose and adjust insulin dosage regimen as needed.

* Interaction does not apply to posaconazole delayed-release tablet.

8 USE IN SPECIFIC POPULATIONS
8.1 Pregnancy

Risk Summary
Published studies, including retrospective cohort studies, national registry studies, and meta-analyses, do not report a consistent pattern or a consistently increased risk of adverse pregnancy-related outcomes with oral use of metoclopramide during pregnancy. However, available data from a case report of GIMOTI use in pregnancy is insufficient to evaluate for a drug-associated risk of major birth defects, miscarriage, or adverse maternal or fetal outcomes.

7 DRUG INTERACTIONS
7.1 Effects of Other Drugs on Metoclopramide

Table 1 displays the effects of other drugs on metoclopramide.

Table 1. Effects of Other Drugs on Metoclopramide

Antipsychotics

Clinical Impact
Potential for additive effects, including increased frequency and severity of tardive dyskinesia (TD), other extrapyramidal symptoms (EPS), and neuroleptic malignant syndrome (NMS).

Intervention
Avoid concomitant use [see Warnings and Precautions (5.1, 5.2, 5.3)].

Strong CYP2D6 Inhibitors, not Included in Antipsychotic Category Above

Clinical Impact
Increased plasma concentrations of metoclopramide; risk of exacerbation of extrapyramidal symptoms [see Clinical

- **Read the step-by-step Instructions for Use that come with your GIMOTI prescription.**
- Take GIMOTI exactly as your healthcare provider tells you. Do not change your dose unless your healthcare provider tells you to.
- GIMOTI comes as a liquid in a glass bottle with a spray pump attached.
- You should not take medicines containing metoclopramide (including GIMOTI) for more than 12 weeks. If you require treatment for longer than 12 weeks, your healthcare provider should frequently monitor you for signs and symptoms of tardive dyskinesia.
- Take GIMOTI at least 30 minutes before each meal and at bedtime.
- If you are not certain that the spray entered the nose, skip the dose and take your next dose at your regularly scheduled time. Do not take an extra dose at the scheduled time. If you miss a dose of GIMOTI, skip that dose and take your next dose at your regular scheduled time. Do not take an extra dose to make up for a missed dose.
- If you take too much GIMOTI, call your poison control center at 1-800-222-1222 or go to the nearest emergency room right away.
- Keep all follow-up visits with your healthcare provider as scheduled, so that your healthcare provider can see how you are responding to treatment with GIMOTI.

What should I avoid while taking GIMOTI?

- Do not drink alcohol while taking GIMOTI. Alcohol may make some side effects of GIMOTI worse, such as feeling sleepy.
- Do not drive, operate machinery, or do other dangerous activities until you know how GIMOTI affects you. GIMOTI may cause sleepiness or dizziness.

- What are the possible side effects of GIMOTI?**
GIMOTI can cause serious side effects, including:
- **Tardive dyskinesia (abnormal muscle movements).** See “What is the most important information I need to know about GIMOTI?”
 - **Other changes in muscle control and movement, such as:**
 - **Uncontrolled spasms of your face and neck muscles, or muscles of your body, arms, and legs (dystonia).** These muscle spasms can cause abnormal movements and body positions and speech problems. These spasms usually start within the first 2 days of treatment. Rarely, these muscle spasms may cause trouble breathing. These spasms happen more often in adults less than 30 years of age and in those who took higher doses of metoclopramide.
 - **Parkinsonism.** Symptoms include slight shaking, body stiffness, and trouble moving or keeping your balance. If you already have Parkinson’s Disease, your symptoms may become worse while you are taking GIMOTI.
 - **Being unable to sit still or feeling you need to move your hands, feet, or body (akathisia).** Symptoms can include feeling jittery, anxious, irritated, or unable to sleep (insomnia), feeling the need to walk around (pacing), and tapping your feet. Your healthcare provider may stop your treatment with GIMOTI if you develop these symptoms.
 - **Neuroleptic Malignant Syndrome (NMS).** NMS is a very rare, but very serious condition that can happen with GIMOTI. NMS can cause death and must be treated in a hospital. Symptoms of NMS include: high fever, stiff muscles, problems thinking, very fast or uneven heartbeat, and increased sweating.
 - **Depression, thoughts about suicide, and suicide.** Some people who take GIMOTI may become depressed, even if they have no history of depression. You may have thoughts about hurting or killing yourself. Some people who have taken metoclopramide have ended their own lives (suicide). Tell your healthcare provider right away if you have these feelings.
 - **High blood pressure.** GIMOTI can cause your blood pressure to increase. Your healthcare provider may stop your treatment if GIMOTI causes your blood pressure to increase very fast.

- **Too much body water.** People who have certain liver problems or heart failure and take GIMOTI may hold too much water in their body (fluid retention). Tell your healthcare provider right away if you have sudden weight gain, or swelling of your hands, legs, or feet.
- **Increased prolactin.** Tell your healthcare provider if your menstrual periods stop or your breasts get larger and make milk. These symptoms go away when you stop taking GIMOTI.

Stop taking GIMOTI, call your healthcare provider, and get medical help right away if you:

- have muscle movements you cannot stop or control
- have muscle movements that are new or unusual
- feel depressed or have thoughts about hurting or killing yourself
- have high fever, stiff muscles, problems thinking, very fast or uneven heartbeat, and increased sweating

The most common side effects of GIMOTI include:

- unpleasant taste
- headache
- tiredness

You may have more side effects the longer you take GIMOTI and the more GIMOTI you take.

You may still have side effects after stopping GIMOTI. You may have symptoms from stopping GIMOTI such as headaches and feeling dizzy or nervous.

Tell your healthcare provider about any side effect that bothers you or that does not go away. These are not all the possible side effects of GIMOTI. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store GIMOTI?

- Store GIMOTI at room temperature between 68°F to 77°F (20°C to 25°C).
- Throw away (discard) GIMOTI 4 weeks after opening even if the bottle contains unused medicine.

Keep GIMOTI and all medicines out of the reach of children.

General information about the safe and effective use of GIMOTI.


Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use GIMOTI for a condition for which it was not prescribed. Do not give GIMOTI to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about GIMOTI that is written for health professionals.

What are the ingredients in GIMOTI?
Active ingredient: metoclopramide
Inactive ingredients: benzalkonium chloride, citric acid monohydrate, edetate disodium dihydrate, purified water, sodium citrate dihydrate, sorbitol.

Manufactured for QOL Medical, LLC by Patheon, a Division of Thermo Fisher. GIMOTI is a trademark of QOL Medical, LLC. ©2026 QOL Medical, LLC. All rights reserved. For more information, go to www.qolmed.com or call 1-833-4-GIMOTI (1-833-444-6684).

This Medication Guide has been approved by the U.S. Food and Drug Administration
Revised: April 2026


- The GIMOTI nasal spray bottle is now ready for use.**
- 3. Using the GIMOTI nasal spray**
- Place the spray nozzle tip under one of your nostrils. Tilt your head slightly forward so the tip of the spray nozzle is aimed away from the center of your nose.



- Close the other nostril with your other index finger. Move the spray pump up so the tip of the nozzle is in your nostril.
-
- Press down firmly on the finger flange until it stops moving to deliver a full dose.
 - Release the finger flange and breathe in gently through the open nostril.
 - Remove the spray pump nozzle tip from your nostril and breathe out slowly through your mouth.
 - After use, wipe the spray nozzle with a clean tissue. Place the cap on the spray nozzle by pushing straight down. Place the safety clip back onto the spray pump.

Cleaning the spray pump nozzle

If the spray pump nozzle becomes clogged, remove it for cleaning by grasping the base of the spray nozzle and pulling up.



Soak the spray nozzle in warm water and rinse. **Do not try** to unblock the spray nozzle by inserting a pin or other sharp object because this will damage the spray nozzle.

Dry the spray nozzle at room temperature. When the spray nozzle is dry, place the dry spray nozzle back on the GIMOTI bottle.

Disposal instructions

The used GIMOTI nasal spray may be thrown away (discarded) in the household trash.

Manufactured for QOL Medical, LLC by Patheon, a Division of Thermo Fisher. GIMOTI is a trademark of QOL Medical, LLC. ©2026 QOL Medical, LLC. All rights reserved.

For more information, go to www.qolmed.com or call 1-833-4-GIMOTI (1-833-444-6684).

These Instructions for Use have been approved by the U.S. Food and Drug Administration
Approved: June 2020

All pregnancies have a background risk of birth defects, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

Clinical Considerations
Fetal/Neonatal Adverse Reactions
Metoclopramide crosses the placental barrier and may cause extrapyramidal signs and methemoglobinemia in neonates with maternal administration during delivery. Monitor neonates for extrapyramidal signs [see *Warnings and Precautions* (5.1, 5.2), *Use in Specific Populations* (8.4)].

Data
Animal Data
Reproduction studies have been performed following administration of oral metoclopramide during organogenesis in pregnant rats at about 6 times the MRHD calculated on body surface area and in pregnant rabbits at about 12 times the MRHD calculated on body surface area. No evidence of adverse developmental effects due to metoclopramide was observed.

8.2 Lactation
Risk Summary
There are no data on the presence of metoclopramide in human milk following nasal administration; however, published data report the presence of metoclopramide in human milk in variable amounts following oral administration [see *Data*]. Systemic exposure following nasal administration of GIMOTI 15 mg is expected to be similar to oral administration of metoclopramide 10 mg [see *Clinical Pharmacology* (12.3)]. Breastfed infants exposed to metoclopramide have experienced gastrointestinal adverse reactions, including intestinal discomfort and increased intestinal gas formation [see *Clinical Considerations*]. Metoclopramide elevates prolactin levels [see *Warnings and Precautions* (5.7)]; however, the published data are not adequate to support drug effects on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother’s clinical need for GIMOTI and any potential adverse effects on the breastfed child from GIMOTI or from the underlying maternal condition.

Clinical Considerations
Monitor breastfeeding neonates because metoclopramide may cause extrapyramidal signs (dystonias) and methemoglobinemia [see *Warnings and Precautions* (5.1, 5.2), *Use in Specific Populations* (8.4)].

Data
In published clinical studies, the estimated amount of metoclopramide received by the breastfed infant was less than 10% of the maternal weight-adjusted oral dose. In one study, the estimated daily amount of metoclopramide received by infants from breast milk ranged from 6 to 24 mcg/kg/day in early puerperium (3 to 9 days postpartum) and from 1 to 13 mcg/kg/day at 8 to 12 weeks postpartum.

8.4 Pediatric Use
Metoclopramide is not recommended for use in pediatric patients due to the risk of tardive dyskinesia (TD) and other extrapyramidal symptoms as well as the risk of methemoglobinemia in neonates. The safety and effectiveness of GIMOTI in pediatric patients have not been established.

Dystonias and other extrapyramidal symptoms associated with metoclopramide are more common in pediatric patients than in adults [see *Indications and Usage* (1), *Warnings and Precautions* (5.1, 5.2)]. In addition, neonates have reduced levels of NADH-cytochrome b₅ reductase, making them more susceptible to methemoglobinemia, a possible adverse reaction of metoclopramide use in neonates [see *Use in Specific Populations* (8.8)].

8.5 Geriatric Use
Metoclopramide is known to be substantially excreted by the kidney, and the risk of adverse reactions, including tardive dyskinesia (TD), may be greater in patients with impaired renal function [see *Warnings and Precautions* (5.1), *Use in Specific Populations* (8.6), *Clinical Pharmacology* (12.3)]. Elderly patients are more likely to have decreased renal function and may be more sensitive to the adverse effects of metoclopramide, especially elderly women, and require a lower starting dosage. GIMOTI is not recommended in geriatric patients as initial therapy. Geriatric patients receiving an alternative metoclopramide product at a stable dosage of 10 mg four times daily can be switched to GIMOTI [see *Dosage and Administration* (2.2)].

8.6 Renal Impairment
The clearance of metoclopramide is decreased and the systemic exposure is increased in patients with moderate to severe renal impairment compared to patients with normal renal function, which may increase the risk of adverse reactions [see *Clinical Pharmacology* (12.3)]. GIMOTI is not recommended in patients with moderate and severe renal impairment (creatinine clearance less than 60 mL/minute), including those receiving hemodialysis and continuous ambulatory peritoneal dialysis [see *Warnings and Precautions* (5.9)]. Use the recommended dosage of GIMOTI in patients with mild renal impairment (creatinine clearance 60 mL/minute or greater) [see *Dosage and Administration* (2.2)].

8.7 Hepatic Impairment
Patients with severe hepatic impairment (Child-Pugh C) have reduced systemic metoclopramide clearance (by approximately 50%) compared to patients with normal hepatic function [see *Clinical Pharmacology* (12.3)]. The resulting increase in metoclopramide blood concentrations increases the risk of adverse reactions. There are no pharmacokinetic data in patients with moderate hepatic impairment (Child-Pugh B). GIMOTI is not recommended in patients with moderate or severe (Child-Pugh B or C) hepatic impairment [see *Warnings and Precautions* (5.9)]. Use the recommended dosage of GIMOTI in patients with mild hepatic impairment (Child-Pugh A) [see *Dosage and Administration* (2.2)].

Metoclopramide, by producing a transient increase in plasma aldosterone, may increase the risk of fluid retention in patients with hepatic impairment [see *Warnings and Precautions* (5.6)]. Monitor patients with hepatic impairment for the occurrence of fluid retention and volume overload.

8.8 NADH-Cytochrome b₅ Reductase Deficiency
Metoclopramide-treated patients with NADH-cytochrome b₅ reductase deficiency are at an increased risk of developing methemoglobinemia and/or sulfhemoglobinemia. For patients with glucose-6-phosphate dehydrogenase (G6PD) deficiency with metoclopramide-induced methemoglobinemia, methylene blue treatment is not recommended. Methylene blue may cause hemolytic anemia in patients with G6PD deficiency, which may be fatal [see *Overdosage* (10)].

8.9 CYP2D6 Poor Metabolizers
Metoclopramide is a substrate of CYP2D6. The elimination of metoclopramide may be slowed in patients who are CYP2D6 poor metabolizers (compared to patients who are CYP2D6 intermediate, extensive, or ultra-rapid metabolizers), possibly increasing the risk of dystonic and other adverse reactions to metoclopramide [see *Clinical Pharmacology* (12.3)]. GIMOTI is not recommended in patients who are CYP2D6 poor metabolizers [see *Warnings and Precautions* (5.9)].

10 OVERDOSAGE
Manifestations of metoclopramide overdosage included drowsiness, disorientation, extrapyramidal reactions, other adverse reactions associated with metoclopramide use (including, e.g., methemoglobinemia), and sometimes death. Neuroleptic malignant syndrome (NMS) has been reported in association with metoclopramide overdose and concomitant treatment with another drug associated with NMS [see *Warnings and Precautions* (5.1, 5.2, 5.3)].

There are no specific antidotes for metoclopramide overdosage. If over-exposure occurs, call your Poison Control Center at 1-800-222-1222 for current information on the management of poisoning or overdosage.

Metoclopramide can be reversed by the intravenous administration of methylene blue. However, methylene blue may cause hemolytic anemia in patients with G6PD deficiency, which may be fatal. Hemodialysis and continuous ambulatory peritoneal dialysis do not remove significant amounts of metoclopramide.

11 DESCRIPTION
Metoclopramide hydrochloride, the active ingredient in GIMOTI, is a dopamine-2 receptor antagonist. Metoclopramide hydrochloride is a white, crystalline, odorless substance, freely soluble in water. Its chemical name is 4-amino-5-chloro-N-[2-(diethylamino)ethyl]-2-methoxy benzamide monohydrochloride monohydrate.

The molecular formula is C₁₄H₂₂ClN₂O₃·HCl·H₂O. Its molecular weight is 354.3. The structural formula is:

GIMOTI (metoclopramide) nasal spray is for nasal administration. The product is supplied as an aqueous solution with a pH of 5.5 ± 0.5 in a 10 mL amber glass vial fitted with a metered spray pump attachment. Each unit contains 9.8 mL.

Each 70 microliter spray contains 15 mg metoclopramide, equivalent to 17.73 mg of metoclopramide hydrochloride. Inactive ingredients consist of benzalkonium chloride, citric acid monohydrate, edetate disodium dihydrate, purified water, sodium citrate dihydrate, and sorbitol.

12 CLINICAL PHARMACOLOGY
12.1 Mechanism of Action
Metoclopramide stimulates motility of the upper gastrointestinal tract without stimulating gastric, biliary, or pancreatic secretions. The exact mechanism of action of metoclopramide in the treatment of gastroesophageal reflux and acute and recurrent diabetic gastroparesis has not been fully established. It seems to sensitize tissues to the action of acetylcholine. The effect of metoclopramide on motility is not dependent on intact vagal innervation, but it can be abolished by anticholinergic drugs.

Metoclopramide increases the tone and amplitude of gastric (especially antral) contractions, relaxes the pyloric sphincter and the duodenal bulb, and increases peristalsis of the duodenum and jejunum resulting in accelerated gastric emptying and intestinal transit. It increases the resting tone of the lower esophageal sphincter. It has little, if any, effect on the motility of the colon or gallbladder.

12.2 Pharmacodynamics
Cardiac Electrophysiology
In a randomized, double-blind, positive-controlled thorough ECG study in 48 healthy subjects, a single administration of 80 mg metoclopramide nasal spray (approximately 5 times the recommended GIMOTI dose) had no effect on the QTc interval.

12.3 Pharmacokinetics
Absorption
The absolute bioavailability of metoclopramide following nasal administration of 10 mg metoclopramide is 47% in healthy subjects compared to intravenous injection of metoclopramide 10 mg. The systemic absorption after nasal administration is lower than that after oral administration given the same dose. Following nasal administration of GIMOTI 15 mg in healthy subjects, the systemic exposure (C_{max} and AUC) to metoclopramide and the time to reach C_{max} (T_{max}) were similar to orally administered 10 mg metoclopramide tablet.

After single nasal administration of metoclopramide at doses ranging from 10 mg to 80 mg in healthy subjects, there was a dose-proportional increase in the mean values for C_{max} and AUC.

The pharmacokinetic parameters of metoclopramide in healthy subjects following a single nasal administration of GIMOTI 15 mg are summarized in Table 3.

Table 3. Summary of Metoclopramide Pharmacokinetic Parameters in Healthy Subjects after a Single Nasal Administration of GIMOTI 15 mg

Parameter ^a	GIMOTI 15 mg
N	94
C _{max} (ng/mL)	41.0 (19.9)
t _{max} (h)	1.25 (0.50 – 3.50)
AUC ₀₋₄ (ng·h/mL)	349 (174.7)
AUC ₀₋₁₂ (ng·h/mL) ^b	367 (184.8)
t _{1/2} (h)	8.1 (2.0)

^aArithmetic mean (SD) except t_{max} for which the median (range) is reported.
^bN = 93

Distribution
Metoclopramide is not extensively bound to plasma proteins (about 30%). The whole body volume of distribution is high (about 3.5 L/kg), which suggests extensive distribution of drug to the tissues.

Elimination
The mean elimination half-life in individuals with normal renal function is approximately 8 hours for administration with GIMOTI 15 mg.

Metabolism: Metoclopramide undergoes enzymatic metabolism via oxidation as well as glucuronide and sulfate conjugation reactions in the liver. Monodethylnmetoclopramide, a major oxidative metabolite, is formed primarily by CYP2D6, an enzyme subject to genetic variability [see *Warnings and Precautions* (5.9), *Use in Specific Populations* (8.9)].

Excretion: Approximately 85% of the radioactivity of an orally administered dose appeared in the urine within 72 hours. After oral administration of 10 or 20 mg, a mean of 18% and 22% of the dose, respectively, was recovered as free metoclopramide in urine within 36 hours.

Specific Populations
Patients with Renal Impairment: In a study of 24 patients with varying degrees of renal impairment (moderate, severe, and end-stage renal disease [ESRD] requiring dialysis), the systemic exposure (AUC) of metoclopramide following oral administration in patients with moderate to severe renal impairment was about 2-fold the AUC in subjects with normal renal function. The AUC of metoclopramide in patients with ESRD on dialysis was about 3.5-fold the AUC in subjects with normal renal function [see *Warnings and Precautions* (5.9), *Use in Specific Populations* (8.6)].

Patients with Hepatic Impairment: In a group of 8 patients with severe hepatic impairment (Child-Pugh C), the average metoclopramide clearance was reduced by approximately 50% compared to patients with normal hepatic function after administration of oral metoclopramide [see *Warnings and Precautions* (5.9), *Use in Specific Populations* (8.7)].

Sex and Body Weight: The AUC₀₋₄ and C_{max} of metoclopramide were 34% and 42% higher in females than in males, respectively, following administration of metoclopramide nasal spray to healthy subjects. Based on population pharmacokinetic analysis, lean body weight (34.3 to 33.5 kg) has a significant impact on metoclopramide pharmacokinetics, with lower systemic exposure expected with higher lean body weight. The clinical significance of these findings is unknown.

Drug Interactions
Effect of Metoclopramide on CYP2D6 Substrates
Although, *in vitro* studies suggest that metoclopramide can inhibit CYP2D6, metoclopramide is unlikely to interact with CYP2D6 substrates *in vivo* at therapeutically relevant concentrations.

Effect of CYP2D6 Inhibitors on Metoclopramide
Metoclopramide 20 mg was orally administered as a single dose to 24 healthy males, without (Period 1) and with (Period 2) a concomitant dose of fluoxetine 60 mg (a strong CYP2D6 inhibitor). Between the two periods, fluoxetine was administered orally for 8 days. The subjects who received concomitant metoclopramide and fluoxetine had a 40% and 90% increase in metoclopramide C_{max} and AUC₀₋₁₂, respectively, compared to subjects who received metoclopramide alone. The mean half-life for metoclopramide was increased from 5.5 (±1.1) hours to 8.5 (±2.2) hours with concomitant fluoxetine [see *Warnings and Precautions* (5.9), *Drug Interactions* (7.1)].

13 NONCLINICAL TOXICOLOGY
13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
Carcinogenesis
A 77-week study was conducted in rats with oral metoclopramide doses up to 40 mg/kg/day (about 6 times the maximum recommended human dose on body surface area basis). Metoclopramide elevated prolactin levels, and the elevation persisted during chronic administration. An increase in mammary neoplasms was found in rodents after chronic administration of metoclopramide [see *Warnings and Precautions* (5.7)]. In a rat model for assessing the tumor promotion potential, a 2-week oral treatment with metoclopramide at a dose of 260 mg/kg/day (about 35 times the MRHD based on body surface area) enhanced the tumorigenic effect of N-nitrosodiethylamine.

Mutagenesis
Metoclopramide was positive in the *in vitro* Chinese hamster lung cell/HGPRT forward mutation assay for mutagenic effects and in the *in vitro* human lymphocyte chromosome aberration assay for clastogenic effects. It was negative in the *in vitro* Ames mutation assay, the *in vitro* unscheduled DNA synthesis assay with rat and human hepatocytes, and the *in vivo* rat micronucleus assay.

Impairment of Fertility
Metoclopramide at intramuscular doses up to 20 mg/kg/day (about 3 times the maximum recommended human dose based on body surface area) was found to have no effect on fertility and reproductive performance of male and female rats.

14 CLINICAL STUDIES
The effectiveness of GIMOTI has been established based on studies of oral metoclopramide for the relief of symptoms in adults with acute and recurrent diabetic gastroparesis.

16 HOW SUPPLIED/STORAGE AND HANDLING
GIMOTI (metoclopramide) nasal spray is supplied as a solution of metoclopramide in a 10 mL Type 1 amber glass bottle fitted with a metered spray pump attachment, a protective cap, and a safety clip. Each box of GIMOTI (NDC 72089-307-15) contains 1 bottle, with FDA-approved Patient Labeling (see Instructions for Use for proper actuation of the device).

Each actuation delivers 15 mg of metoclopramide. Each bottle contains 9.8 mL which is sufficient for 4 weeks of 4 times a day use.

Store at 20°C to 25°C (68°F to 77°F) excursions permitted 15°C to 30°C (59°F and 86°F) [see *USP Controlled Room Temperature*].

Discard GIMOTI 4 weeks after opening even if the bottle contains unused medicine.

17 PATIENT COUNSELING INFORMATION
Advise the patient or caregiver to read the FDA-approved patient labeling (Medication Guide and Instructions for Use).

Adverse Reactions
Tardive Dyskinesia and/or other Extrapyramidal Reactions
Inform patients that GIMOTI may cause tardive dyskinesia or other extrapyramidal symptoms, parkinsonian symptoms, and motor restlessness. Instruct patients to immediately discontinue GIMOTI and contact their healthcare provider if symptoms occur [see *Warnings and Precautions* (5.1, 5.2)].

Neuroleptic Malignant Syndrome
Inform patients that serious neuroleptic malignant syndrome (NMS) has been reported in association with concomitant treatment with another drug associated with NMS. Advise patients to report all prescription and over-the-counter medications to the healthcare provider. Instruct patients to immediately discontinue GIMOTI and seek medical attention if symptoms occur [see *Warnings and Precautions* (5.3)].

Depression and/or Possible Suicidal Ideation
Inform patients that symptoms of new onset or worsening depression as well as suicidal ideation have been reported in patients taking metoclopramide. Instruct patients to immediately discontinue GIMOTI and contact their healthcare provider if any of these symptoms occur [see *Warnings and Precautions* (5.4)].

Drug Interactions
Distribution
Inform the patients or their caregivers that concomitant treatment with numerous other medications can precipitate or worsen serious adverse reactions such as tardive dyskinesia or other extrapyramidal reactions, neuroleptic malignant syndrome, and CNS depression [see *Drug Interactions* (7.1, 7.2)]. Explain that the prescriber of any other medication must be made aware that the patient is taking GIMOTI.

Effects on the Ability to Drive and Operate Machinery
Inform the patient or their caregiver that GIMOTI may cause drowsiness or dizziness, or otherwise impair the mental and/or physical abilities required for the performance of hazardous tasks such as operating machinery or driving a motor vehicle [see *Warnings and Precautions* (5.8)].

Administration Instructions [see *Dosage and Administration* (2.1)]
Advise the patients or their caregiver to read the *Instructions for Use* on how to appropriately administer GIMOTI:

- One spray in one nostril administers the appropriate dose.
- Before administering the first dose from a bottle, prime the pump by pressing down on the finger flange and releasing 10 sprays in the air.
- Place the spray nozzle tip under one nostril and lean the head slightly forward so the tip of spray nozzle is aimed away from the septum and toward the back of the nose.
- Close the other nostril with the other index finger. Move spray pump upwards so the tip of the nozzle is in the nostril.
- To ensure a full dose, hold the bottle upright while pressing down firmly and completely on finger flange and release while inhaling slowly through the open nostril.
- Remove spray pump nozzle tip from nostril and exhale slowly through the mouth.
- Wipe the spray nozzle with a clean tissue.

Missed or Incomplete Doses

- If uncertain that the spray entered the nose, do not repeat the dose. Take the next dose at the scheduled time.
- If a dose is missed, take the next dose of GIMOTI at the regularly scheduled time. Do not make up for the missed dose or double the next dose.

Manufactured for: QOL Medical, LLC
Vero Beach, FL 32963 USA
By: Patheon, a Division of Thermo Fisher
Bourgain Jallieu Cedex, 38307, France
© 2026 QOL Medical, LLC. All rights reserved.